

Second virial coefficient for a d -dimensional Lennard-Jones ($2n$ - n) system

M.L. Glasser

Department of Physics and Clarkson Center for Statistical Physics
Clarkson University
Potsdam, NY 13699-5820 (USA)

This note examines the second virial coefficient for an imperfect gas subject to a $2n$ - n interparticle potential in any dimension d between 0 and n . A compact analytic expression is presented for this quantity which shows that, apart from a numerical factor, its temperature dependence is a universal function parameterized by d/n .

PACS:05.70.Ce, 02.30.+g

Keywords: Lennard-Jones potential, Virial coefficient.

The virial expansion has played an important role in studying imperfect gases and has recently been invoked for the examination of systems, such as alkanes absorbed at interfaces [1], in two dimensions. It is therefore of some interest to look at the dimensional dependence of the coefficients. In this note an analytic expression is presented for the second virial coefficient in the case of the Lennard-Jones 2n-n potential

$$V(r) = 4\epsilon[(\frac{\sigma}{r})^{2n} - (\frac{\sigma}{r})^n] \quad (1)$$

for spatial dimension $0 < d < n$, where it is defined.

The second virial coefficient is given by

$$B_d(T) = \frac{1}{2} \int d^d r [1 - \exp(-V(r)/kT)]. \quad (2)$$

Since $V(r)$ is spherically symmetric, the angular integrals in hyperspherical coordinates can be carried out to give

$$B_d(T) = \frac{\pi^{d/2}}{\Gamma(d/2)} \int_0^\infty r^{d-1} [1 - e^{-V(r)/kT}] dr. \quad (3)$$

After scaling out σ , integration by parts, and an elementary change of integration variable, (3) reduces to

$$B_d(T) = C_d [2I_{1-d/6} - I_{-d/6}] \quad (4)$$

where

$$\begin{aligned} a &= 4\epsilon/kT \\ C_d &= \frac{\pi^{d/2} \sigma^d}{d\Gamma(d/2)} \\ I_\nu &= a \int_0^\infty x^\nu e^{-a(x^2-x)} dx. \end{aligned} \quad (5)$$

The latter integral does not appear to have been tabulated, but is easily evaluated as follows: Replace $\exp(ax)$ by its Taylor series and integrate term-by-term using Euler's gamma function integral to get an infinite series summed with respect to n , say. Then break this up into even ($n = 2k$) and odd ($n = 2k + 1$) terms. In these series replace $n!$ by $4^k k! (1/2)_k$ and $4^k k! (3/2)_k$,

respectively. The resulting series are seen to be Kummer functions[2] and we obtain

$$I_\nu = \frac{1}{2}a^{-(\nu-1)/2}[\Gamma(\frac{\nu+1}{2})M(\frac{\nu+1}{2}, \frac{1}{2}, \frac{a}{4}) + \sqrt{a}\Gamma(\frac{\nu}{2}+1)M(\frac{\nu}{2}+1, \frac{3}{2}, \frac{a}{4})]. \quad (6)$$

Therefore, for $0 < d < n$, in terms of the dimensionless temperature $\tau = 4/a = kT/\epsilon$, and $\rho = d/2n$

$$\begin{aligned} B_{d,n}(\tau) = & -\frac{\sigma^d \pi^{d/2} 2^{d/n}}{2n\Gamma(d/2)} \tau^{-\rho} [\Gamma(-\rho)M(-\rho, \frac{1}{2}, \tau^{-1}) + \\ & 2\tau^{-1/2}\Gamma(\frac{1}{2}-\rho)M(\frac{1}{2}-\rho, \frac{3}{2}, \tau^{-1})] = \\ & \frac{\sigma^d \pi^{d/2} 2^{2\rho}}{n\Gamma(d/2)} \tau^{-\rho} e^{1/\tau} [\Gamma(-\rho)M(\rho + \frac{1}{2}, \frac{1}{2}, -1/\tau) + \\ & \frac{2}{\sqrt{\tau}}\Gamma(\frac{1}{2}-\rho)M(\rho + 1, \frac{3}{2}, -1/\tau)] = \\ & \frac{\sigma^d \pi^{(d-1)/2} 2^{2\rho}}{n\Gamma(d/2)} \Gamma(-\rho)\Gamma(\frac{1}{2}-\rho)D_{2\rho}(-\sqrt{2/\tau}). \end{aligned} \quad (7)$$

In deriving (7) the identities

$$\begin{aligned} M(1 - \frac{d}{2n}, \frac{1}{2}, z) - 2zM(1 - \frac{d}{2n}, \frac{3}{2}, z) &= M(-\frac{d}{2n}, \frac{1}{2}, z) \\ (1 - \frac{d}{n})M(\frac{3}{2} - \frac{d}{2n}, \frac{3}{2}, z) - M(\frac{1}{2} - \frac{d}{2n}, \frac{1}{2}, z) &= -\frac{d}{n}M(\frac{1}{2} - \frac{d}{2n}, \frac{3}{2}, z) \end{aligned} \quad (8)$$

and Kummer's first identity have been used. The last expression in (7) is particularly attractive, but as indicated below, the parabolic cylinder function can be problematic and the expressions using Kummer functions are probably better for numerical work. From Eq.(7) we have the interesting fact that, apart from a constant prefactor, the scaled temperature dependence of the second virial coefficient is a universal function of d/n . Thus, the Boyle temperature (at which the second virial coefficient vanishes), is given by the zeros of the parabolic cylinder function. Listed below are the Boyle temperatures for the L-J 12-6 potential in various dimensions:

d	τ_B	d	τ_B
0.5	0.45812	3.5	5.35369
1	0.72701	4	9.01642
1.5	1.07421	4.5	17.1531
2	1.56031	5	41.0597
2.5	2.28098	5.5	173.8854
3	3.41793		

The Boyle temperature is therefore also universal with respect to d/n , so this table lists the Boyle temperatures for $d/n = 1/12, \dots, 11/12$.

Finally, we present the Joule-Thompson coefficient $J_{d,n}(\tau) = \tau \partial B_{d,n}(\tau) / \partial \tau - B_{d,n}(\tau)$.

$$J_{d,n}(\tau) = \frac{\sigma^d \pi^{d/2} 2^{d/2n}}{6n\Gamma(d/2)} \tau^{-d/2n} \{ 3\Gamma(-\frac{d}{2n}) [(1 + \frac{d}{2n}) M(-\frac{d}{2n}, \frac{1}{2}, \tau^{-1}) - \frac{2}{\tau} \frac{d}{2n} M(1 - \frac{d}{2n}, \frac{3}{2}, \tau^{-1})] + 2\tau^{-1/2} \Gamma(\frac{1}{2} - \frac{d}{2n}) [3(\frac{3}{2} + \frac{d}{2n}) M(\frac{1}{2} - \frac{d}{2n}, \frac{3}{2}, \tau^{-1}) + \frac{2}{\tau} (\frac{1}{2} - \frac{d}{2n}) M(\frac{3}{2} - \frac{d}{2n}, \frac{5}{2}, \tau^{-1})] \}. \quad (12)$$

A similar calculation for the case $d = 3$ was carried out a number of years ago by Garrett [4] who expressed his result in terms of a sum of parabolic cylinder functions $D_\nu(-z)$. This function appears to be an orphan among special functions, in that there is very little information about it in standard references such as [3]. What there is concerns the function $D_\nu(z)$. In spite of the similarity in notation, the two are independent solutions of the parabolic cylinder equation and their relationship is not a simple one. For example, for real z ,

$$D_{1/2}(z) = \frac{1}{\sqrt{\pi}} (z/2)^{3/2} [K_{\frac{1}{4}}(z^2/4) + K_{\frac{3}{4}}(z^2/4)] \quad (13)$$

while

$$D_{1/2}(-z) = \frac{\sqrt{\pi}}{4} z^{3/2} [I_{\frac{3}{4}}(z^2/4) + I_{-\frac{3}{4}}(z^2/4) - I_{\frac{1}{4}}(z^2/4) - I_{-\frac{1}{4}}(z^2/4)]. \quad (14)$$

On the other hand, the Kummer function used in (7) (also denoted $M(a, b, c) = {}_1F_1(a; b; c)$) is standard in mathematical packages and does not suffer from

this problem. For $d = 3$ (7) is consistent with Garrett's formula. Recently, Vargas et al.[5] have treated $B_{3,6}$ by an improvement of Garrett's approach and have expressed the virial coefficient in terms of modified Bessel functions of fractional order. By making use of (13) and (14), (7) can be transformed into their expression. Finally, the method used here for the $2n - n$ potential: series expansion, decomposition of the sum into relatively prime summands and the use of the Gauss multiplication formula, can be applied to the general $m - n$ potential and leads to a closed form expression as a sum of $lcm(m, n)/n$ hypergeometric functions. Thus, for the $12 - 5$ potential considered in [1], the second virial coefficient was expressed as a sum of 12 Kummer functions, which proved to be of little practical value.

References

- [1]. B.A. Pethica, M.L. Glasser and J. Mingins, J. Colloid and Interface Sci. **81**, 41 (1981).
- [2]. A. Lou and B.A. Pethica, Langmuir **13**, 4933 (1997).
- [3]. **Handbook of Mathematical Functions**, [Eds. M. Abramowitz and I. Stegun, NBS Applied Math. Ser. 55, Washington (1964)].
- [4]. A.J.M. Garrett, J.Phys. A:Math. Gen., **13**, 379 (1980).
- [5]. P. Vargas, E. Muños and L. Rodriguez, Physica A **290**, 92 (2001).